## illumına<sup>®</sup>

# STM Reagent Improves Robustness of Infinium® Methylation Assay

#### Introduction

The Infinium Methylation Assay detects cytosine methylation (an important factor in many human diseases) based on highly multiplexed genotyping of bisulfite-converted genomic DNA (gDNA). Upon treatment with bisulfite, unmethylated cytosines are deaminated to uracil, while methylated cytosines remain unaffected. The assay interrogates these chemically-differentiated loci using two site-specific probes, one designed for the methylated locus and another for the unmethylated. Single-base extension of the probes incorporates a DNP-or biotin-labeled ddNTP, which is subsequently stained with a fluorescence reagent in order to determine the signal intensity ratio of methylated vs unmethylated residues<sup>1</sup>.

The X-Stain protocol used for fluorescence staining has been updated with an improved reagent that delivers highly robust performance across HumanMethylation27 BeadChips. According to the conventional X-Stain protocol, the hybridized probes are single-base extended, then stained using Labeling Two-Color Master Mix (LTM)<sup>2</sup>. In the updated protocol, LTM is replaced by Superior Two-Color Master Mix (STM), the standard reagent for all Infinium HD BeadChips.

#### Analysis of Data Quality and Consistency

To evaluate the consistency of the data produced by each reagent, replication experiments were performed with six samples that were independently assayed using either STM or LTM. The correlation coefficients for all probe signals indicated a very strong linear relationship between replicates. Evaluation of the individual samples showed  $r^2 > 0.99$  in all cases tested (Table 1). Overall, this analysis demonstrated that STM and LTM produce highly consistent data, meaning that samples processed with either stain can be analyzed within the same GenomeStudio<sup>®</sup> project.

A comparison study was performed over a larger range of samples to evaluate how the reagents performed across a more diverse sample set. Forty-eight reference gDNA samples were assayed using both STM and LTM to compare the CpG detection accuracy. As illustrated in Figure 1, STM showed reduced variability and an increased percentage of known CpG sites detected from the reference samples. These results suggest that STM provides more robust data quality and better reproducibility over a broader range of samples and BeadChips for the Infinium Methylation Assay.

### **Updated Staining Procedure**

All Infinium Assays based on HD BeadChips use STM as the standard staining reagent. Although the HumanMethylation27 BeadChip is not HD-formatted, STM can be incorporated into the assay protocol with only minor procedural modifications.

For the automated assay protocol, an update to the Tecan robot setup is recommended. In the updated procedure, once the BeadChips have been placed in the Chamber Rack and slid into the robot bed, the XStain HD BeadChip task should be selected on the robot PC (Figure 2). After selecting this task, the STM reagent tubes can be placed on the robot in the same location that the LTM tubes would have occupied. After this point, the assay can be completed as normal by following the conventional protocol.

For the manual assay protocol, the only required procedural change is to replace the LTM reagent tubes with STM tubes before starting an experiment.



The above table shows correlations between beta values, which represent methylation levels of the CpG sites. Analysis of replicates treated with STM and LTM show a strong correlation, with r2 > 0.99 across all samples tested.



#### Conclusions

Replacing LTM with STM in the Infinium Methylation Assay produces results that are highly consistent under either reagent condition. However, when evaluated over a broader range of samples and Bead-Chips, the STM reagent proves to be more robust, and can modestly improve the signal detection accuracy of the Infinium Methylation Assay.

Figure 2: Tecan Robot Setup Modification for the Assat Automated Protocol



To prepare the Tecan robot for STM, select XStain Tasks | Infinium II Chemistry | X-Stain HD BeadChip

#### Additional Information

Visit our website or contact us at the address below to learn more about Infinium products and protocols.

Illumina, Inc. • 9885 Towne Centre Drive, San Diego, CA 92121 USA • 1.800.809.4566 toll-free • 1.858.202.4566 tel • techsupport@illumina.com • illumina.com

#### FOR RESEARCH USE ONLY

© 2010 Illumina, Inc. All rights reserved.

Illumina, illumina, Dx, Solexa, Making Sense Out of Life, Oligator, Sentrix, GoldenGate, GoldenGate Indexing, DASL, BeadArray, Array of Arrays, Infinium, BeadXpress, VeraCode, IntelliHyb, iSelect, CSPro, GenomeStudio, Genetic Energy, HiSeq, and HiScan are registered trademarks or trademarks of Illumina, Inc. All other brands and names contained herein are the property of their respective owners. Pub. No. 270-2009-001 Current as of 10 Aug 2009

